



Chemical Responses of *Nicotiana tabacum* (Solanaceae) Induced by Vibrational Signals of a Generalist Herbivore

Carlos F. Pinto¹ · D. Torrico-Bazoberry¹ · M. Penna² · R. Cossio-Rodríguez¹ · R. Cocroft³ · H. Appel⁴ · H. M. Niemeyer¹

Received: 27 October 2018 / Revised: 1 July 2019 / Accepted: 7 July 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

Abstract

Plants are able to sense their environment and respond appropriately to different stimuli. Vibrational signals (VS) are one of the most widespread yet understudied ways of communication between organisms. Recent research into the perception of VS by plants showed that they are ecologically meaningful signals involved in different interactions of plants with biotic and abiotic agents. We studied changes in the concentration of alkaloids in tobacco plants induced by VS produced by *Phthorimaea operculella* (Lepidoptera: Gelechiidae), a generalist caterpillar that naturally feeds on the plant. We measured the concentration of nicotine, normicotine, anabasine and anatabine in four treatments applied to 11-weeks old tobacco plant: a) Co = undamaged plants, b) Eq = Playback equipment attached to the plant without VS, c) Ca = Plants attacked by *P. operculella* herbivory and d) Pl = playback of VS of *P. operculella* feeding on tobacco. We found that nicotine, the most abundant alkaloid, increased more than 2.6 times in the Ca and Pl treatments as compared with the Co and Eq treatments, which were similar between them. Normicotine, anabasine and anatabine were mutually correlated and showed similar concentration patterns, being higher in the Eq treatment. Results are discussed in terms of the adaptive significance of plant responses to ecologically important VS stimuli.

Keywords Nicotine · Plant-insect interactions · Tobacco · Plant response · Plant perception · Plant signaling · Herbivore

Introduction

Studies on behavior and memory have traditionally been performed on animals since it has been assumed that such traits are absent in plants (Applewhite 1975; Trewavas 2017; Vertosick 2002). Nevertheless, recent research has shown that plants can sense their biotic and abiotic environments and selectively respond to many types of stimuli. A number of functions have been described in plants which are analogous to those in animals but differ in mechanisms and capabilities (Karban 2015). For example, plants can assess the external environment, deal with predators and diseases, exhibit flexible reproductive strategies, all in the absence of a central nervous

system (Trewavas 2017). Legg and Hutter (2007) propose that intelligence in organisms depends on the ability of individuals to adapt to different objectives and environments and allows them to interact with the environment and succeed when exposed to different challenges. Plants can fit with the previous description, since they exhibit phenotypic and molecular responses to environmental signals, and they are able to improve their fitness through behavioral changes which eventually become adaptive strategies during their life cycle (Karban 2015; McNamara and Houston 1996; Trewavas 2017).

Plants may be exposed to a variety of stimuli (e.g. light, chemicals, wind, temperature, sounds) (Karban 2015; Telewsky 2006) and they are able to scan and sense the environment and respond to prevailing conditions (Karban 2015; Mischra et al. 2016). Light sensing involves phytochromes, cryptochromes and phototropins; chemical sensing involves chemoreceptors; mechanical sensing is performed by mechanoreceptors; and it is not yet clear which specific receptors are responsible for temperature sensing (Karban 2015; Wagner et al. 2008). Although mechanisms of sound or vibrational sensing are just starting to be disentangled, a body of knowledge about their perception and the related feedback by plants has been produced (Appel and Cocroft 2014; Braam 2005; Helms et al. 2013; Karban 2015; Smith 2000; Wagner et al. 2008).

✉ H. M. Niemeyer
niemeyer@abulafia.ciencias.uchile.cl

¹ Facultad de Ciencias, Universidad de Chile, Santiago, Chile

² Programa de Fisiología y Biofísica, Facultad de Medicina, Universidad de Chile, Santiago, Chile

³ Division of Biological Sciences, University of Missouri, Columbia, MO 65211, USA

⁴ Department of Environmental Sciences, University of Toledo, Toledo, OH 43606, USA

Vibrational signals (VS) constitute one of the oldest ways of communication between organisms, with an estimated number of 230,000 species of invertebrates and many vertebrates using them (Cocroft et al. 2014; Hill and Wessel 2016). In spite of their widespread occurrence, however, VS remain one of the most understudied communication modes, far behind other modalities such as chemical and visual signals (Cocroft et al. 2014). Recent research has made important advances in the perception of VS by plants and their ecological consequences (Appel and Cocroft 2014; Body et al. 2019; De Luca and Vallejo-Marin 2013; Gagliano et al. 2012; Gagliano 2013, Gagliano et al. 2018). When plants perceive VS they can modify or adjust processes such as root growth, seed germination, defense responses, and responses to water scarcity (Appel and Cocroft 2014; Body et al. 2019; Chowdhury et al. 2014; Gagliano 2013; Jeong et al. 2014). Hence, VS have been recognized as meaningful signals involved in the interaction of plants with their environment (Mischra et al. 2016).

In the realm of plant–insect interactions, the chemical arsenal of plants against herbivory has been studied at different levels and scales (Baluška and Ninkovic 2010), including constitutive internal defensive secondary metabolites, chemicals induced by herbivore attack, and organic volatile compounds that repel herbivores, attract natural enemies of herbivores or alert other plant parts and individuals (Kessler and Baldwin 2002; Meiners 2015; Meinwald and Eisner 2008). Several species of the genus *Nicotiana* (Solanaceae) have been extensively studied from all these approaches, and nicotine and related alkaloids have been described as constitutive and induced defensive metabolites (Baldwin 1988, 1989; Halitschke et al. 2000).

The ecological importance of VS for plants is highlighted by work on the defensive responses of *Arabidopsis thaliana* (Brassicaceae) to feeding vibrations of a specialist predator, *Pieris rapae* (Pieridae). Caterpillar feeding vibrations, alone or in concert with simulated herbivory, can activate hormonal signaling pathways (Body et al. 2019), prime glucosinolate and phenolic based leaf defenses (Appel and Cocroft 2014), and modulate the release of volatile organic compounds from leaves (Body et al. 2019). However, the role of herbivore feeding vibrations in plant defense responses has not been documented in other plant species. Here, we investigated changes in alkaloid concentration in *Nicotiana tabacum* L. induced by VS produced by *Phthorimaea operculella* (Lepidoptera: Gelechiidae), a generalist caterpillar which naturally feeds in the plant leaves.

Materials and Methods

Plants *Nicotiana tabacum* (Solanaceae) plants were grown in a greenhouse at the Universidad de Chile. Lighting was provided by metal halide lamps (36 W) and the room was kept at $25 \pm$

2°C with a 16:8 h L:D photoperiod. Seeds were germinated in peat (Kekkilä; Vantaa, Finland) and after six weeks of development seedlings were transplanted into pots (14-cm tall \times 17-cm diameter) containing a mix of potting soil (Armony; Pudahuel, Chile), sand (Armony; Pudahuel, Chile) and peat in a 4:1:1 proportion. Plants were watered twice weekly, 1 day with water and the other with an aqueous fertilizer solution (Phostrogen Plant Food, Santiago, Chile; <http://www.bayergarden.co.uk/Products/p/Phostrogen-All-Purpose-Plant-Food>).

Insect Rearing and Caterpillar Herbivory Patterns

Phthorimaea operculella, commonly named tobacco splitworm or potato tuberworm, is a global pest of solanaceous crops and weeds (Rondon 2010). Individuals were collected near Curacaví (Metropolitan Region, Chile: 33.49°S , 71.02°W , 185 m above sea level) and reared on *N. tabacum* under the same laboratory conditions as the plants. *P. operculella* caterpillars are leaf-miners, feeding mostly from the mesoderm of leaves (Rondon 2010). Adults of *P. operculella* were maintained in cages containing 3-month old tobacco plants. Adults were fed with 10% aqueous honey solution every 2 days (Golizadeh et al. 2014), while caterpillars were allowed to feed *ad libitum* on the tobacco plants until the fourth instar, which was the stage used for bioassays.

Natural patterns of caterpillar feeding were characterized using 20 4th-instar caterpillars. Individuals were placed on the fifth leaf of a tobacco plant (counted from the apex of the plant) and observed for 8 h to characterize: a) temporal patterns of feeding and b) leaf area attacked over time. Photographs of individuals and leaves were taken every 15 min and were analyzed using the software MorphoJ. The amount of leaf area attacked over time constituted the baseline for designing Herbivory and Playback bioassays.

Feeding Vibration Recordings and Playback Setup

Vibrational signals produced by a feeding 4th-instar *P. operculella* caterpillar were recorded using the same conditions as in insect rearing. The caterpillar was placed inside a clip cage on the fifth leaf of an 11-week old tobacco plant ($N=20$), once the caterpillar started to feed, the clip cage was removed. Vibrations were recorded for 10 min per individual with a laser Doppler vibrometer (Polytec CLV-2534) positioned at less than 1 cm from where the caterpillar was feeding.

To reproduce the vibrations produced by the feeding caterpillar, a 9 mm Samsung/Mplus Linear Resonant Actuator (LRA) vibrator was used, driven by a Behringer HA8000 V2 headphone amplifier. An accelerometer (Vibrametrics 9002A with P5000 power supply) was attached with wax to the LRA to calibrate the playback. The LRA and accelerometer were attached to the leaf using a modified hair clip covered with EVA rubber, held in position with a helping hands tool (Quad Hands). The clip held the actuator lightly but securely against the leaf surface; our previous observations

indicate that this procedure avoided injury to the leaf. Before each playback bioassay, the frequency response and amplitude of the actuator were characterized after attaching it to the leaf. Because an accelerometer was used to calibrate playback of laser-recorded signals, the playback recordings were converted from velocity to acceleration using Matlab. Digital filters were applied using the signal processing toolbox of MATLAB R2017a to compensate for the frequency response of the playback setup, and produce playbacks that closely matched the temporal and spectral properties of the original recordings (Appel and Cocroft 2014; Cocroft 2010; Cocroft et al. 2014).

The playback design was based on the natural pattern of feeding behavior of *P. operculella* caterpillars (see Results). Caterpillar vibrations were played back to the fifth leaf of naive tobacco plants ($n = 18$ replicates). Three feeding vibration exemplars with a high signal-to-noise ratio were used, each from a different individual of *P. operculella* feeding on a different tobacco plant.

Plant Chemical Defenses Alkaloids were extracted based on a modification of the protocol by Saitoh et al. (1985). The fifth leaf of each plant used during bioassays was cut, immediately flash frozen with liquid nitrogen and pulverized in a porcelain mortar. The fifth leaf was selected because it exhibits greater induced alkaloidal responses on the fifth day after stimulation (Baldwin 1989, 1999). The powdered leaf was transferred to a glass vial and weighed. Five mL of methanol were added and the vial was shaken at 600 rpm for 30 min. After that, the methanolic extract was filtered through filter paper (Whatman No. 1) and transferred into a Florence flask where it was evaporated to dryness using a rotary evaporator at 40 °C. After adding 5 mL of chloroform to the Florence flask, it was immersed in a 25 °C ultrasound bath for 10 min. The extract was then transferred into a separatory funnel and 5 mL of 5% hydrochloric acid were added, thoroughly mixed and the organic phase collected and discarded; the acidic extract was washed twice with 5 mL of chloroform. After that, 5 mL of 29% aqueous ammonia solution was added and the basic solution was extracted twice with 5 mL of chloroform. The organic phases were collected in a vial and then evaporated to dryness under a nitrogen flow.

The alkaloidal extract was redissolved in a mixture of 70 μ L of chloroform and 30 μ L of internal standard solution (0.25 mM docosane in chloroform). Then, 1 μ L of the redissolved extracts was injected in a Shimadzu model GCMS-QP 2010 Ultra gas chromatograph (Shimadzu, Kyoto, Japan). The mass spectrometer used electron impact (EI) ionization mode (70 eV) with an emission current of 250 μ A. The temperatures of the injection port, transfer line and ion source were 250 °C, 280 °C and 250 °C, respectively. An Rtx-5MS Crossbond 5% diphenyl 95% dimethyl polysiloxane capillary GC column (30 m length, 0.25 mm I.D., 0.25 μ m film thickness) (Restek, Bellefonte, PA, USA)

was used in the splitless mode, with helium as the carrier gas at 50 ml/min. The GC oven temperature was programmed to remain at 30 °C for 3 min, then to increase to 230 °C at a rate of 18 °C/min and finally to remain at 230 °C for 5 min. Retention indexes were determined based on chromatograms obtained from the periodic injection of a standard alkane mixture (Sigma-Aldrich). Compounds were identified based on comparisons of their retention index and mass spectrum with those in the NIST14 database and with authentic standards.

To quantify the four main alkaloids (nicotine, nor nicotine, anabasine and anatabine) present in the extracts, calibration curves were constructed for each alkaloid with five concentrations of commercial standards of each one (Sigma-Aldrich) ranging from 0.0224 to 2 mg/mL and containing the same concentration of internal standard described above (Yuan et al. 2018). The concentration of each alkaloid in the extracts was extrapolated from the corresponding calibration curve, and was expressed as μ g/g of fresh sample.

Bioassays In all bioassays, 11-week old plants were used. Four types of bioassays were performed. In all of them, alkaloids were quantified in the fifth leaf of the tobacco plants which had been subjected to the following treatments: a) Constitutive, corresponding to the concentration in the undamaged leaf ($N = 18$) (Kaplan et al. 2008). b) Caterpillar, characterizing the chemical response of tobacco to herbivory by *P. operculella* caterpillars. One 4th instar caterpillar was placed on the fifth leaf of an undamaged plant, and enclosed within a clip cage ($N = 18$) in order to limit its movements before it started to feed. The individual was observed every 10 min to determine the start of feeding. Once the caterpillar started to feed, the clip cage was removed, and the caterpillar allowed to eat ad-libitum for 208 min. After this time, it was carefully removed from the plant and the plant was left in the rearing environment described above. Five days after caterpillar removal, the damaged leaf was removed, and alkaloids were extracted and quantified. c) Playback. The same protocol was used as in the Caterpillar treatment except that the caterpillar was replaced by the recording of VS produced by an individual caterpillar chewing the leaf, as described above ($N = 18$). d) Equipment, characterizing the potential effect of the setup used for playback on the induction of secondary metabolites. The complete playback setup was turned on and attached to the fifth leaf of the plant without any sound being reproduced for the same time as the playback replicates ($N = 18$) in order to evaluate the effect of the equipment by itself (e.g. weight, electrical fields). After 5 days the leaf was collected and chemically analyzed. The experimental treatments were conducted during the same time period, with 3–4 plants treated per day, each with a different treatment.

Statistical Analyses A Principal Component Analysis (PCA) was applied to all replicates of alkaloid concentrations in all treatments on R 3.4.4 (R Core Team 2014). Thereafter,

treatments were compared within each alkaloid using General Linear Models (GLM) also on R 3.4.4 (R Core Team 2014). Linear regression was used to generate calibration curves for the alkaloid standards. Pearson correlations were used to evaluate potential associations between the concentrations of different alkaloids.

Results

Caterpillars fed in bouts of 9.70 ± 1.35 min (mean \pm sd) with a frequency of 2.48 ± 0.26 bouts per hour and a pause (no feeding) lasting 15.98 ± 1.85 min between feeding events. The waveform of chewing vibrations reflected the repetitive nature of the chewing process (Fig. 1). These data were used in the design of the playback stimulus, which consisted of 1 min of the original feeding vibrations recordings from one individual repeated 10 times (i.e. 10-min stimulus) followed by a 16-min silent pause. This basic 26-min pattern was repeated 8 times, so, the complete stimulus lasted for 208 min. Based on the




Fig. 1 Chewing vibrations of *P. operculella* on *N. tabacum*. **a** Waveform; **b** Spectrogram; **c** Amplitude spectrum

caterpillar observations described above, during this time period an average of 14.4 ± 1.5 (mean \pm sd) % of the leaf area was attacked by a single caterpillar.

The coefficients of determination (R^2) for calibration curves were 0.94 for nicotine, 0.96 for normicotine, 0.98 for anabasine and 0.95 for anatabine. The most abundant alkaloid in all treatments was nicotine, followed by anatabine, normicotine and anabasine (mean concentrations over all treatments: nicotine = $873 \mu\text{g/g}$, anatabine = $216.4 \mu\text{g/g}$, normicotine = $77.3 \mu\text{g/g}$, anabasine = $33.7 \mu\text{g/g}$) (Fig. 2).

The PCA with all the alkaloid concentration data showed that most of the variability (94%) was explained with the first two components, the first one involving normicotine, anabasine and anatabine (74% of variability) and the second one involving only nicotine (21% of variability). The corresponding biplot showed two subgroups, one where Playback and Caterpillar treatments overlapped and another with Equipment and Constitutive treatments overlapped (Fig. 2). Nicotine levels differed between treatments (Table 1), being 1.7 times higher in plants of the Equipment treatment, 2.9 times higher in plants of the Caterpillar treatment and 2.8 times higher in plants of the Playback treatment, with respect to the Constitutive treatment.




Fig. 2 Biplot of the PCA applied to alkaloid concentrations (*Nic* Nicotine, *Nor* Normicotine, *An* Anabasine and *Anat* Anatabine) of the four treatments: *Co* Constitutive, *Eq* Equipment, *Ca* Caterpillar, *Pl* Playback

Table 1 Results for the Generalized Linear Model for the concentrations of alkaloids expressed as $\mu\text{g/g}$ of fresh sample. Treatments were: *Co* Constitutive, *Eq* Equipment, *Ca* Caterpillar and *Pl* Playback (see text for description of treatments)

	Estimate	Standard error	t value	Pr(> z)
a) Nicotine: ANOVA: $F = 3.8326$ $p = 0.01367^*$ AIC: 1145.8				
Co vs Eq	94.4	329.6	0.286	0.7755
Co vs Ca	814.1	307.8	2.645	0.0102
Co vs Pl	789.6	307.8	2.565	0.0126
Pl vs Eq	-695.2	333.5	-2.085	0.0410
Pl vs Ca	24.6	311.9	0.079	0.9375
Eq vs Ca	719.7	333.5	2.158	0.0346
b) Nornicotine: ANOVA: $F = 6.4086$ $p < 0.001$ AIC: 786.27				
Co vs Eq	85.99	26.77	3.212	0.0021
Co vs Ca	-21.95	25.04	-0.876	0.3841
Co vs Pl	-10.50	25.04	-0.419	0.6765
Pl vs Eq	96.49	26.77	3.604	0.0006
Pl vs Ca	-11.45	25.04	-0.457	0.6490
Eq vs Ca	-85.99	26.77	-3.212	0.0021
c) Anabasine: ANOVA: $F = 8.1874$ $p < 0.001$ AIC: 615.5				
Co vs Eq	24.149	7.627	3.166	0.0024
Co vs Ca	-1.640	7.135	-0.230	0.8189
Co vs Pl	-13.156	7.135	-1.844	0.0698
Pl vs Eq	37.305	7.627	4.891	7.08e-06
Pl vs Ca	11.516	7.135	1.614	0.1114
Eq vs Ca	-25.789	7.627	-3.381	0.00123
d) Anatabine: ANOVA: $F = 3.4086$ $p = 0.02261$ * AIC: 946.06				
Co vs Eq	174.34	77.51	2.249	0.0279
Co vs Ca	-66.48	72.38	-0.918	0.3618
Co vs Pl	-14.88	72.38	-0.206	0.8377
Pl vs Eq	189.22	78.42	2.413	0.0186
Pl vs Ca	-51.60	73.36	-0.703	0.4843
Eq vs Ca	-240.82	78.42	-3.071	0.0031

The GLM analysis showed that the concentration of nicotine in the Playback and Caterpillar treatments did not differ significantly and both differed from those of the Constitutive and Equipment treatments, which had similar nicotine levels (Table 1, Fig. 3). Nornicotine, anabasine and anatabine concentrations differed significantly between bioassays, being higher and significantly different in the Equipment treatment compared with the other three treatments (Table 1, Fig. 3).

Concentrations of nornicotine were highly and positively correlated with concentrations of anabasine ($R = 0.85$, $p < 0.001$) and of anatabine ($R = 0.95$, $p < 0.001$). Furthermore, concentrations of anabasine were positively correlated with concentrations of anatabine ($R = 0.83$, $p < 0.001$).

Discussion

Plants of *N. tabacum* chemically responded to natural herbivory by *P. operculella* caterpillars and to VS produced by feeding activity of *P. operculella*. Increased concentration of nicotine was observed under two situations: when leaves were damaged by a caterpillar, and when leaves were stimulated with playbacks of caterpillar chewing vibrations. On the other hand, nicotine concentration did not increase when the equipment for playback was attached to the plant and turned on.

Nicotine constitutes the first effective defensive barrier in the genus *Nicotiana* because it is the most abundant alkaloid in the plant and also because it can be further induced in response to herbivore attack and physical damage (Baldwin 1996, 1999; Baldwin et al. 1997; Halitschke et al. 2000; Stepphun et al. 2004). In fact, plants can sense even small scratches caused by the walking of an insect on the leaf surface and as a consequence activate nicotine induction (Peiffer et al. 2009; Karban 2015). The present study demonstrates for the first time in *N. tabacum* that isolated vibrations produced by a feeding caterpillar can induce a chemical response by the plant. This result shows that tobacco plants can perceive this stimulus, recognize it as a potential threat and choose to invest energy in the induction of defenses. On the other hand, plants do not respond chemically in the same way to different types of damage (e.g. different herbivore species, diseases). Herbivores can alter the ability to affect chemical defenses and even suppress their induction following damage (Tooker et al. 2008; Alba et al. 2011; Witzany and Baluška 2012). For example, in the genus *Nicotiana*, the specialist moth *Manduca sexta* can disrupt the biosynthesis of nicotine in the roots through their oral regurgitant (McCloud and Baldwin 1997; Kahl et al. 2000; Halitschke et al. 2000, 2004). The present




Fig. 3 Concentration of alkaloids expressed as $\mu\text{g/g}$ of fresh sample for each treatment. Different letters (a, b) indicate significant differences between treatments at $P < 0.05$. Treatments: *Co* Constitutive, *Eq* Equipment, *Ca* Caterpillar, *Pl* Playback

study leaves open the question of which features of the played-back feeding vibrations were important for triggering the defensive responses, and of what responses may be evoked by vibrations produced by other herbivores or other biotic or abiotic sources. Phenolic defenses of *Arabidopsis* plants were induced by playback of *Pieris* feeding vibrations, but not by playback of wind vibrations or leafhopper song (Appel and Cocroft 2014), revealing some specificity in the vibration-induced defensive response. However, the key features of herbivore vibrations necessary for inducing plant responses, and the range of variation in effective vibrational stimuli, have so far not been identified.

The minor alkaloids, nornicotine, anabasine and anatabine were found at significantly higher levels in the Equipment treatment as compared with all other treatments, consistent with the correlations found between these variables. However, the mechanisms leading to these concentration increases are not clear. We lack an explanation for the observation that the Playback treatment did not lead to an increase in the concentration of any of the minor alkaloids. Attaching the equipment should have had the same positive effect in both the 'Equipment' and 'Playback' treatments, and it is possible that the vibrational stimulus downregulated the production of the minor alkaloids.

Plant response to herbivory can be expressed in a variety of traits (e.g. morphology, chemistry, development) (Karban 2015; Karban and Baldwin 1997) and induced chemical responses by the plant may cause herbivores to be less attracted to the defended tissue or may reduce the performance of herbivores that have consumed such tissue (Karban 2015; Karban and Myers 1989). Our experimental approach has assessed plant induced chemical responses to VS isolated from the other effects of insect feeding, such as the mechanical damage to the plant and the insect's saliva, thereby providing a deeper understanding of the role of VS in plant responses to herbivory. We suggest that VS of herbivory constitute an ecological stimulus to *N. tabacum* and is the main reason for nicotine induction as a behavioral response of the plant interprets such signals as a threat. These results, taken together with similar observations made on *A. thaliana* (Appel and Cocroft 2014) suggest that plant perception and responses to VS might be widespread in plants. The adaptive value of the chemical response to VS remains unclear: the inclusion of fitness traits into the experimental design will be necessary to assess these responses as adaptive behaviors expressed by plants.

Acknowledgements We thank CONICYT (Comisión Nacional de Ciencia y Tecnología) postdoctoral project 3160356, LANBIO (Latin American Network for Research on Bioactive Natural Compounds), ISP (International Science Programs at Uppsala University (project BOL-01), and National Science Foundation IOS-1359593 for financial and logistic support. We are also indebted to David Veliz for his input in the statistical analyses and Ian T. Baldwin from the Max Planck Institute for Chemical Ecology for providing a sample of anatabine.

Compliance with Ethical Standards

Conflict of Interest The authors declare that they have no conflict of interest.

References

- Alba JM, Glas JJ, Schimmel BCJ, Kant MR (2011) Avoidance and suppression of plant defenses by herbivores and pathogens. *J Plant Interact* 6:221–227. <https://doi.org/10.1080/17429145.2010.551670>
- Appel HM, Cocroft RB (2014) Plants respond to leaf vibrations caused by insect herbivore chewing. *Oecologia* 175:1257–1266. <https://doi.org/10.1007/s00442-014-2995-6>
- Applewhite PB (1975) Plant and Animal Behavior: An Introductory Comparison. In: Eisenstein EM (ed) *Aneural Organisms in Neurobiology*. Advances in Behavioral Biology, vol 13. Springer, Boston, pp 131–139
- Baldwin IT (1988) Damage-induced alkaloids in tobacco: pot-bound plants are not inducible. *J Chem Ecol* 14:1113–1120. <https://doi.org/10.1007/BF01019339>
- Baldwin IT (1989) Mechanism of damage-induced alkaloid production in wild tobacco. *J Chem Ecol* 15:1661–1680. <https://doi.org/10.1007/BF01012392>
- Baldwin IT (1996) Methyl jasmonate-induced nicotine production in *Nicotiana attenuata*: inducing defenses in the field without wounding. *Entomol Exp Appl* 80:213–220. https://doi.org/10.1007/978-94-009-1720-0_49
- Baldwin IT (1999) Inducible nicotine production in native *Nicotiana* as an example of adaptive phenotypic plasticity. *J Chem Ecol* 25:3–30. <https://doi.org/10.1023/A:1020880931488>
- Baldwin IT, Zhang Z-P, Diab N, Ohnmeiss TE, McCloud ES, Lynds GY, Schmelz EA (1997) Quantification, correlations and manipulations of wound-induced changes in jasmonic acid and nicotine in *Nicotiana sylvestris*. *Planta* 201:397–404. <https://doi.org/10.1007/s004250050082>
- Baluška F, Ninkovic V (2010) *Plant communication from an ecological perspective*. Springer, Uppsala
- Body MJA, Neer WC, Vore C, Lin CH, Vu DC, Schultz JC, Cocroft RB, Appel HM (2019) Caterpillar chewing vibrations cause changes in plant hormones and volatile emissions in *Arabidopsis thaliana*. *Front Plant Sci* 10 (810)
- Braam J (2005) In touch: plant responses to mechanical stimuli. *New Phytol* 165:373–389. <https://doi.org/10.1111/j.1469-8137.2004.01263.x>
- Chowdhury MEK, Lim H, Bae H (2014) Update on the effects of sound wave on plants. *Res Plant Dis* 20:1–7
- Cocroft RB (2010) Vibrational communication. In: Breed M & J Moore (eds.) *Encyclopedia of animal behavior*. Elsevier, Amsterdam. Pp 498–505
- Cocroft RB, Gogala M, Hill PSM, Wessel A (2014) Studying vibrational communication. In: Janik VM, McGregor P (eds) *Animal Signals and Communication*. Springer, Berlin, pp 1–61
- De Luca PA, Vallejo-Marin M (2013) What's the 'buzz' about? The ecology and evolutionary significance of buzz-pollination. *Curr Opin Plant Biol* 16:429–435. <https://doi.org/10.1016/j.pbi.2013.05.002>
- Gagliano M (2013) Green symphonies: a call for studies on acoustic communication in plants. *Behav Ecol* 24:789–796. <https://doi.org/10.1093/beheco/ars206>
- Gagliano M, Renton M, Duvdevani N, Timmins M, Mancuso S (2012) Out of sight but not out of mind: alternative means of communication in plants. *PLOS ONE*. <https://doi.org/10.1371/journal.pone.0037382>

- Gagliano M, Abramson CI, Depczynski M (2018) Plants learn and remember: lets get used to it. *Oecologia* 186:29–31. <https://doi.org/10.1007/s00442-017-4029-7>
- Golizadeh A, Esmaceli N, Razmjou J, Rafiee-Dastjerdi H (2014) Comparative life tables of the potato tuberworm, *Phthorimaea operculella*, on leaves and tubers of different potato cultivars. *J Insect Sci* 14:42. <https://doi.org/10.1093/jis/14.1.42>
- Halitschke R, Kessler A, Johannes K, Lorenz A, Baldwin IT (2000) Ecophysiological comparison of direct and indirect defenses in *Nicotiana attenuata*. *Oecologia* 124:408–417. <https://doi.org/10.1007/s004420000389>
- Halitschke R, Kessler A, Kahl J, Lorenz A, Baldwin IT, Dudareva N, Pichersky E, Gershenzon J (2004) Biochemistry of plant volatiles. *Plant Physiol* 135:1893–1902. <https://doi.org/10.1104/pp.104.049981>
- Helms AM, De Moraes CM, Tooker JF, Mescher MC (2013) Exposure of *Solidago altissima* plants to volatile emissions of an insect antagonist (*Eurosta solidaginis*) deters subsequent herbivory. *Proc Natl Acad Sci USA* 110:199–204. <https://doi.org/10.1073/pnas.1218606110>
- Hill P, Wessel A (2016) Biotremology. *Curr Biol* 26(5):187–191. <https://doi.org/10.1016/j.cub.2016.01.054>
- Jeong MJ, Cho JI, Park SH, Kim KH, Lee SK, Kwon TR, Park SC, Siddiqui ZS (2014) Sound frequencies induce drought tolerance in rice plant. *Pak J Bot* 46:2015–2020
- Kahl J, Siemens DH, Aerts RJ, Gabler R, Kuhnemann F, Preston CA, Baldwin IT (2000) Herbivore-induced ethylene suppresses a direct defense but not a putative indirect defense against an adapted herbivore. *Planta* 210:336–342. <https://doi.org/10.1007/PL00008142>
- Kaplan I, Halitschke R, Kessler A, Sardanelli S, Denno RF (2008) Constitutive and induced defenses to herbivory in above- and belowground plant tissues. *Ecology* 89:392–406
- Karban R (2015) *Plant Sensing and Communication*. University of Chicago Press, Chicago
- Karban R, Baldwin IT (1997) *Induced responses to herbivory*. University of Chicago Press, Chicago
- Karban R, Myers J (1989) Induced plant responses to herbivory. *Annu Rev Ecol Syst* 20:331–348
- Kessler A, Baldwin IT (2002) Plant responses to insect herbivory: The emerging molecular analysis. *Annu Rev Plant Biol* 53:299–328. <https://doi.org/10.1146/annurev.arplant.53.100301.135207>
- Legg S, Hutter M (2007) A collection of definitions of intelligence. FAIA 157:17–24 <https://arxiv.org/abs/0706.3639v1>. Accessed Oct 2018
- McCloud ES, Baldwin IT (1997) Herbivory and caterpillar regurgitants amplify the wound-induced increases in jasmonic acid but not nicotine in *Nicotiana sylvestris*. *Planta* 203:430–435. <https://doi.org/10.1007/s004250050210>
- McNamara JM, Houston AI (1996) State dependent life histories. *Nature* 380:215–221
- Meiners T (2015) Chemical ecology and evolution of plant-insect interactions: A multitrophic perspective. *Curr Opin Insect Sci* 8:22–28. <https://doi.org/10.1016/j.cois.2015.02.003>
- Meinwald J, Eisner T (2008) Chemical Ecology in retrospect and prospect. *Proc Natl Acad Sci USA* 105:4539–4540. <https://doi.org/10.1073/pnas.0800649105>
- Mischra RC, Ghosh R, Bae H (2016) Plant acoustics in the search of a sound mechanism for sound signalling in plants. *J Exp Bot* 67:4483–4494. <https://doi.org/10.1093/jxb/erw235>
- Peiffer M, Tooker JF, Luthe DS, Felton GW (2009) Plants on early alert: glandular trichomes as sensors for insect herbivores. *New Phytol* 184:644–656. <https://doi.org/10.1111/j.1469-8137.2009.03002.x>
- Rondon SI (2010) The potato tuberworm: a literature review of its biology, ecology, and control. *Am J Potato Res* 87: 149–166
- Saitoh F, Noma M, Kawashima N (1985) The alkaloid contents of 60 *Nicotiana* species. *Phytochemistry* 24: 4774–480
- Smith H (2000) Phytochromes and light signal perception by plants –an emerging synthesis. *Nature*. 407:585–591. <https://doi.org/10.1038/35036500>
- Stephuna A, Gase K, Krock B, Halitschke R, Baldwin IT (2004) Nicotine's defensive function in nature. *PLOS Biol* 2(10):e382. <https://doi.org/10.1371/journal.pbio.0020382>
- Telewsky FW (2006) A unified hypothesis of mechanoreception in plants. *Am J Bot* 93:1466–1476. <https://doi.org/10.3732/ajb.93.10.1466>
- Tooker JF, Rohr JR, Abrahamson WG, De Moraes CM (2008) Gall insects can avoid and alter indirect plant defenses. *New Phytol* 178: 657–671. <https://doi.org/10.1111/j.1469-8137.2008.02392.x>
- Trewavas A (2017) The foundations of plant intelligence. *Interface Focus* 7:20160098. <https://doi.org/10.1098/rsfs.2016.0098>
- Vertosick FT (2002) *The Genius Within. Discovering the Intelligence of Every Living Thing*. Harcourt Inc, New York
- Wagner AM, Krab S, Wagner MJ, Moore AL (2008) Regulation of thermogenesis in flowering *Araceae*: the role of the alternative oxidase. *Biochim Biophys Acta* 1777:993–1000. <https://doi.org/10.1016/j.bbabi.2008.04.001>
- Witzany G, Baluška F (2012) *Biocommunication of Plants*. Springer, Berlin
- Yuan Y, Zhou R, Li DL, Luo C, Li GY (2018) Simultaneous quantitative assessment of nine glycosides in tobacco by liquid chromatography-tandem mass spectrometry. *J Sep Sci* 41:1009–1016. <https://doi.org/10.1002/jssc.201700880>